

CATALOGUE  
OF THE  
L I Z A R D S

IN THE  
BRITISH MUSEUM  
(NATURAL HISTORY).

---

SECOND EDITION.

BY  
GEORGE ALBERT BOULENGER.

VOLUME III.  
LACERTIDÆ, GERRHOSAURIDÆ, SCINCIDÆ,  
ANELYTROPIDÆ, DIBAMIDÆ,  
CHAMÆLEONTIDÆ.

LONDON:  
PRINTED BY ORDER OF THE TRUSTEES.  
1887.

LIBRARY

Femoral pores 16 to 21. Tail about twice as long as head and body; upper caudal scales truncate, strongly keeled. Olive-grey above, back uniform or scantily dotted with black; a blackish band or series of black spots on each side, extending to the eye or the loreal region; sides with round white spots; lower surfaces uniform greenish white (in spirit).

	♂. millim.	♀. millim.
Total length .....	232	193
Head .....	21	15
Width of head .....	15	10
From end of snout to fore limb.....	32	25
From end of snout to vent.....	80	71
Fore limb .....	28	24
Hind limb .....	46	39
Tail .....	152	122

## Syria.

<i>a.</i> ♀.	— ?	[Type.]
<i>b-d.</i> ♂ ♀.	Damascus.	Dr. J. Anderson [P.].
<i>e-g.</i> ♂ ♀.	Jerusalem.	Canon Tristram [C.].
<i>h.</i> ♂.	Dead Sea.	Canon Tristram [C.].

18. *Lacerta jayakari*. (PLATE II.)

Head moderately depressed; snout long, obtusely acuminate. Rostral entering the nostril; two superposed postnasals; six upper labials anterior to the subocular; a series of granules between the supraoculars and the supraciliaries; occipital broader than the interparietal; temple minutely and uniformly granulate, the granules much smaller than the dorsal scales; tympanic shield very small; a large anterior supratemporal shield, followed by a series of smaller ones. No trace of a gular fold; 47 to 50 gular scales on a line between the collar and the third pair of chin-shields; collar with very slightly serrated edge, composed of 13 or 15 small plates. Dorsal scales very small, granular, round, convex, smooth or very indistinctly keeled, 85 to 95 across the middle of the body; four or five lateral scales correspond to the length of the ventral shield. Ventrals broader than long, in eight longitudinal series, the outer series narrowest, the others equal; 27 or 28 transverse series. Preanal small, surrounded by two or three semicircles of smaller plates. The hind limb reaches the collar. Scales on upper surface of crus slightly larger than dorsals, obtusely keeled. A series of large shields under the arm. Femoral pores 26 to 28. Tail about twice as long as head and body; upper caudal scales truncate, strongly keeled. Olive above; adult uniform, white-spotted on the cheeks and sides of the neck; young spotted with black and whitish all over the body, with three black vertical bars on the temple and a black spot on each labial shield. Lower surfaces uniform yellowish.

	millim.		millim.
Total length .....	472	From end of snout to	
Head .....	38	vent.....	152
Width of head .....	27	Fore limb .....	56
From end of snout to		Hind limb.....	90
fore limb .....	65	Tail .....	320

Arabia.

a-b. ♂ & yg. Near Muscat. A. S. G. Jayakar, Esq. [P.].

### 19. *Lacerta galloti*.

*Zootoca derbiana*, Gray, *Cat.* p. 29.

— *galloti*, part., Gray, *l. c.* p. 30.

*Lacerta galloti*, Dum. & Bibr. v. p. 238; Boettg. *Abh. Senckenb. Ges.* ix. 1874, p. 174; Bedriaga, *Arch. f. Nat.* 1879, p. 325; Bouleng. *Proc. Zool. Soc.* 1881, p. 740; Bedriaga, *Abh. Senckenb. Ges.* xiv. 1886, p. 301.

Head moderately depressed; snout long, obtuse. Rostral not entering the nostril; a single postnasal; five upper labials anterior to the subocular; a series of granules between the supraoculars and the supraciliaries; occipital, in the adult, longer and broader than the interparietal; temporal scales granular; masseteric and tympanic shields distinct; no large supratemporals. Gular fold indicated; 40 to 46 gular scales on a line between the collar and the third pair of chin-shields; collar with even or very slightly serrated edge, composed of 9 to 13 small plates. Dorsal scales very small, roundish-rhomboidal, juxtaposed, obtusely keeled, 84 to 96 across the middle of the body; three or four lateral scales correspond to the length of a ventral shield. Ventrals in 12 or 14 longitudinal and 29 to 31 transverse series. Præanal small, bordered by two or three semicircles of small plates. The hind limb reaches the shoulder or the collar. Upper crural scales smaller than dorsals. Femoral pores 23 to 31. Tail at least twice as long as head and body; upper caudal scales truncate, strongly keeled. Olive-grey or dark olive above, with black and pale greenish spots; females and young with two light, black-edged longitudinal streaks on each side. Lower surfaces olive or blackish in the adult, paler in the young; throat, in the young, with angular grey bands converging forwards. A more or less distinct black band and a few whitish spots on the hinder side of the thighs.

	♂. millim.	♀. millim.
Total length .....	305	277
Head .....	30	28
Width of head .....	24	19
From end of snout to fore limb ....	52	48
From end of snout to vent .....	115	112

Pl. II.



Mintert. Eros. imp.

*Lacerta jayakari*

BRIT. MUS. N.H.

2/2

P. Smit del et lith.