

- Lacerta viridis*, *Boettger*, *Zeitschr. Ges. Naturw.* (3) iv. 1879, p. 505.  
 ———, var. *gadovii*, *Bouleng. Proc. Zool. Soc.* 1884, p. 438,  
 pl. xxxviii.; *Bedriaga*, *Abh. Senckenb. Ges.* xiv. 1886, p. 74.  
 ———, var. *schreiberi*, *Bedriaga*, *l. c.* p. 76.

Approaching *L. ocellata*. Granules between the supraoculars and the supraciliaries usually reduced to three or four; occipital rather large, usually broader than the interparietal; no distinct tympanic shield. Dorsal scales smaller, less distinctly hexagonal, less strongly keeled than in the typical form; 52 to 58 scales across the middle of the body; lateral scales larger than median dorsals. Usually 8 well-developed rows of ventrals. Femoral pores 12 to 17, usually 13 or 14. Young olive or brownish above, with large yellow or bluish-white, black-edged ocelli on the sides of the head and body. These ocelli often disappear in the adult, which are green or brown above, variously spotted or reticulated with black. Lower surfaces yellowish, immaculate in the young, often with round black spots in the adult. Throat blue in the adult.

	♂. millim.	♀. millim.
Total length.....	260	287
Head .....	23	22
Width of head.....	16	15
From end of snout to fore limb ..	36	36
From end of snout to vent .....	90	97
Fore limb.....	30	33
Hind limb .....	46	47
Tail .....	170	190

Spain and Portugal.

- |                                       |                              |  |
|---------------------------------------|------------------------------|--|
| <i>a-f.</i> ♂ & yg.                   | Coruña.                      | M. V. L. Seoane [P.].                                  |
| <i>g-i, k-m.</i> ♂, ♀,<br>hgr., & yg. | Serra de Gerez.              | Dr. H. Gadow [C.].                                     |
| <i>n.</i> ♀.                          | Coimbra.                     | Dr. H. Gadow [C.].                                     |
| <i>o-r.</i> ♂, ♀, & yg.               | Serra de Monchique, Algarve. | Dr. H. Gadow [C.].<br>(Types of var. <i>gadovii</i> .) |

C. Var. *major*.

- Lacerta viridis*, *Bibr. & Bory de St. Vinc. Expéd. Sc. Morée, Zool.*  
 p. 66, pl. x. fig. 1; *Bedriaga*, *Bull. Soc. Nat. Mosc.* 1881, p. 74.  
 ——— *quinquevittata* (*non Mén.*), *Erhard, Faun. Cycl.* i. p. 80.  
*Tropidosaura algira* (*non L.*), *Erhard, l. c.*  
*Lacerta viridis*, vars. *punctata* et *trilineata*, part., *Bedriaga*, *Abh.*  
*Senckenb. Ges. t. c.*

A larger form, attaining nearly to the size of *L. ocellata*. Constantly a series of granules between the supraoculars and the supraciliaries; occipital usually as broad as or a little broader but shorter than the interparietal; temporal scales smaller than in the typical form usually with a very distinct tympanic shield. 50 to 58 scales across the middle of the body, the laterals not larger than the

median dorsals. Usually a well-developed additional series of ventrals, making in all eight longitudinal rows. Femoral pores 15 to 20. Young olive above, usually with three or five yellowish longitudinal streaks, the lowermost of which, extending from axilla to groin, is often replaced by a series of round spots. These bands usually disappearing in the adult, which are green on the body and limbs, uniform or finely speckled with black; upper surface of head vermiculated with black; lower surfaces yellow, the outer ventrals often speckled with black; throat never blue.

	♂. millim.	♀. millim.
Total length.....	415	350
Head .....	36	26
Width of head .....	27	17
From end of snout to fore limb ..	62	42
From end of snout to vent .....	155	115
Fore limb.....	49	37
Hind limb .....	83	61
Tail .....	260	235

Dalmatia, Turkey, Greece, Asia Minor.

a. ♂.	Dalmatia.	Dr. Heckel.
b-c. ♂.	Greece.	
d. ♀.	Syra (?).	
e-f. ♂ & hgr.	Rhodes.	C. G. Danford, Esq. [P.].
g, h-m. ♀ & yg.	Crete.	Baron v. Maltzan [C.].
n. ♂.	Asia Minor.	C. Fellows, Esq. [P.].
o-r. Yg.	Smyrna.	
s. ♂.	Zebil Bulgar Dagh, Cili- cian Taurus, 4000 feet.	C. G. Danford, Esq. [P.].

#### D. Var. *strigata*.

*Lacerta strigata*, Gray, *Cat.* p. 32.

*Lacerta strigata*, *Eichw. Zool. Spec. Ross. Pol.* iii. p. 189, and *Faun.*

*Casp.-Cauc.* p. 70, pl. x. figs. 4-6; *Anders. Proc. Zool. Soc.* 1872, p. 372; *Blanf. Zool. E. Pers.* p. 364.

— *quinquevittata*, *Ménétr. Cat. Rais.* p. 61.

— *viridis*, *Kessler, Tr. St. Petersb. Nat. Soc.* viii. 1878, p. 146;

*Boettg. Ber. Senckenb. Ges.* 1879-80, p. 170, and in *Radde, Faun. Flor. Casp. Geb.* p. 39.

— —, var. *strigata*, *Bedriaga, Abh. Senckenb. Ges. t. c.* p. 91.

Size moderate. A series of granules between the supraoculars and the supraciliaries; occipital small, much shorter than the interparietal; tympanic shield distinct. 40 to 48 scales across the middle of the body; dorsals strongly keeled. Only six longitudinal rows of ventrals. Femoral pores 16 to 20. Young greenish or olive, spotted or marbled with black, and with three or five light dorsal streaks, which may disappear completely in the adult; these may be closely dotted with black, as certain specimens of the