

Sites 7 and 8: Rio Negro province: 25 de Mayo department: 7) Ruta Provincial 5, 40 km SE Maquinchao (41° 30' 07.2" S, 68° 33' 26.3" W, 887 m). 28 November 2000. L. J. AVILA, C. H. F. PEREZ. LJAMM 3089 (female). 8) Ruta Nacional 23, 14 km W Aguada de Guerra (41° 00' 37.2" S, 68° 30' 23.0" W, 866 m). 28 November 2000. L. J. AVILA, C. H. F. PEREZ. LJAMM 3090 (female).

All Rio Negro province specimens are females and although identification is difficult we assign these individuals to *P. nigroiugulus* according to the description of CEI et al. (2001). In site 7, we found the lizard below a small rock in a place characterized by rocky basaltic outcrops with vegetation characteristic of the shrubby-grass steppe of the Patagonian Steppe formation with *Mulinum*, *Stipia*, *Senecio*, and *Adesmia* as dominant plants. Other lizard species observed in sympatry were *Liolaemus bibronii*, *L. cf. boulengeri* KOSLOWSKY, 1898, *L. cf. melanops*, *L. kriegi* MÜLLER & HELLMICH, 1939, *L. petrophilus*, *Homonota darwinii*, *Diplolaemus darwinii*, *Phymaturus patagonicus*, and *Leiosaurus belli* DUMÉRIL & BIBRON, 1837. In site 8, landscape was a plain with consolidated soil, with a plant and lizard community similar to site 7; the lizard was collected in late afternoon basking on a shrub branch. Citations of CRUZ et al. (1999) of *Pristidactylus scapulatus* (BURMEISTER, 1861) for 77 km NW Sierra Colorado (2 km S La Esperanza) on the roadside of Provincial Road 68, Department 25 de Mayo, as well as the individual mentioned for Chubut province, probably will have to be assigned to *P. nigroiugulus*, however, we did not yet revise these individuals.

REFERENCES: AVILA, L. J. (1994): Geographic Distribution: *Pristidactylus achalensis*.- Herpetol. Rev., Hays; 25: 165-166. AVILA, L. J., & MONTERO, R. & MORANDO, M. (2000): Evaluación del estado de conservación de la fauna de lagartijas y anfisbénidos de la República Argentina; 108 pp. In: LAVILLA, E. O. & RICHARD, E. & SCROCCHI, G. J. (Eds.): Categorización de anfibios y reptiles de Argentina. Tucuman (Asoc. Herpetol. Argentina). AVILA, L. J. & MORANDO, M. & PEREZ, D. R. (2001): New records and natural history notes for lizards and snakes from Patagonia, Argentina.- Herpetol. Rev., Hays; 32: 64-65. CEI, J. M. (1986): Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas áridas y semiáridas.- Mus. Reg. Sci. Nat., Torino; Monografie IV, 527 pp. CEI, J. M. (1993). Reptiles del noroeste, nordeste y este de la Argentina. Herpetofauna de las selvas subtropicales, Puna y Pampas.- Mus. Reg. Sci. Nat.,

Torino; Monografie XI., 929 pp. CEI, J. M. & SCOLARO, J. A. & VIDELA, F. (2001): The present status of Argentinean polychrotid species of the genus *Pristidactylus* and description of its southernmost taxon as a new species.- J. Herpetol., Athens; 35: 597-605. CRUZ, F. B., & SCHULTE II, J. A. & BELLAGAMBA, P. (1999): New distributional records and natural history notes for reptiles from southern Argentina.- Herpetol. Rev., Hays; 30: 182-183. ETHERIDGE, R. & ESPINOZA, R. E. (1997): New records and natural history notes for lizards from north-western Argentina.- Herpetol. Rev., Hays; 28: 160-161. LAMBOROT, M. & DIAZ, N. F. (1987): A new species of *Pristidactylus* (Sauria: Iguanidae) from central Chile and comments on the speciation in the genus.- J. Herpetol., Athens; 21:29-37.

KEY WORDS: Reptilia: Squamata: Sauria: Iguanidae: *Pristidactylus nigroiugulus*, chorology, distribution, Río Negro and Chubut provinces, Argentina

SUBMITTED: April 15, 2003

AUTHORS: Luciano Javier AVILA, Mariana MORANDO, CONICET-CRILAR, Mendoza and Entre Rios s/n, 5301, Anillaco (La Rioja), Argentina. (Present address: Department of Integrative Biology, Brigham Young University, 401 WIDB, Provo, Utah, USA) < luciano.javier@hotmail.com >; Cristian Hernan Fulvio PEREZ, Agustín Alvarez 1182 "A", 8000. Bahía Blanca (Buenos Aires), Argentina. < liolaemu@criba.edu.ar >

Ectoparasite of lizards and possible vector: the mammal hard tick *Haemaphysalis concinna* KOCH, 1844

During a herpetological excursion to the Croatian island of Krk in the middle of May 2002, ten tick-infested lizards [*Algyroides nigropunctatus* (DUMÉRIL & BIBRON, 1839), *Podarcis melisellensis* (BRAUN, 1877), *P. muralis* (LAURENTI, 1768), *Lacerta bilineata* DAUDIN, 1802], and a tick-infested snake [*Elaphe longissima* (LAURENTI, 1768)] were detected. The reptilian hosts originating from two localities (44°58,943 N / 14°27,79 E; 47 m a.s.l.; 45°06,088 N / 14°20,99 E; 120 m a.s.l.; description of the flora and herpetofauna see SEHNAL 1999 and RATHBAUER 2002) were freed from their pests on site. The ticks were killed and fixed in 70% ethanol. Blood drops emerging on this occasion were smeared on glass slides and air dried for later staining according to Giemsa.

Altogether 46 ticks were collected (table 1), 35 could be determined to species level and, according to the key of BABOS (1964), turned out to be unfed nymphs of *Haemaphysalis concinna* KOCH, 1844 (Ixodidae, Acari) as is undoubtedly demon-

Table 1: Number of tick nymphs found attached to eleven specimens of reptiles on the island of Krk (Croatia). * - blood parasites detected.

Reptile taxon	n (reptiles)	n (nymphs)
<i>Podarcis muralis</i>	2	8 / 3*
<i>Podarcis melisellensis</i>	6	6*/5/4*/3/1*/1*
<i>Lacerta bilineata</i>	1	11
<i>Algyroides nigropunctatus</i>	1	1
<i>Elaphe longissima</i>	1	3
Σ	11	46

strated by the shape of the scutum and the palps (fig. 1). Unfortunately, the number of hosts was too low to generate epidemiological data.

Nymphs of *Haemaphysalis sulcata* (CANESTRINI & FANZAGO, 1877) are long known reptilian parasites in Asia (HOOGSTRAAL et al. 1981), but this species has never been found in Central Europe; and Central European lizards have never been found infested with *H. concinna* (e.g., REHACEK et al. 1961). *Haemaphysalis concinna* is a triannual, three-host species with minor host preference. Common hosts are larger mammals like cattle, sheep and goat, red deer and deer, and sometimes even man (MERDIVENCI 1969). The most frequent tick parasitizing lizards in Europe, *Ixodes ricinus* (LINNEUS, 1758) (JANSEN 2002), is a species which is able to suck blood from its reptilian victims without perceptible harm, even if they do so in very high numbers

Fig. 1: Nymph of *Haemaphysalis concinna* from a lizard. The picture combines a photo taken in visual light, an auto-fluorescence photo (scutum), and a line drawing. Note the shape of the palps, of the scutum, and the festoons. Tick length: 1.4 mm.




(MALKMUS 1995). Nothing is known about the pathogenic potential of *Haemaphysalis* spp. to reptiles; yet the presence of unfed ticks only may be interpreted as a hint for the existence of a poor adaptation of this tick species to the reptilian hosts. Nevertheless, except for the unusual host spectrum, all other ecological parameters of our tick finding, like activity season, species distribution area, and microclimate, were within the typical range (NOSEK et al. 1967). The area of tick infestation on the lizards' bodies was restricted to the trunk close to the insertion of the anterior limbs. This may be due to the ability of lizards to detach ticks by feeding and by scratching with the limbs.

The killing of ticks by feeding on them is the main infection route of lizards with blood parasites of the genus *Hepatozoon* (Adeleina, Apicomplexa). *Haemaphysalis* ticks are known vectors or intermediate, maybe even final hosts of this blood parasite. Thus, we looked for infections of the reptiles with this intraerythrocytic parasites by screening stained blood smears. Five cases of *Hepatozoon* infections were detected (table 1, fig. 2). Unfortunately, without knowing the vegetative stages within the reptile's organs, usually the liver, a species determination of this parasite is impossible.

Our observations of *H. concinna* ticks being attached to reptiles are remarkable for two reasons: (i) Not one fully engorged tick has been found, only more or less unfed

Fig. 2: Blood smear, Giemsa stained, from *Podarcis muralis* showing an intraerythrocytic parasite, most probably a gametocyte of a tick transmitted *Hepatozoon* species. Bar represents 10 µm.



nymphs. Thus, reptile hosts may be inexpensive for this tick species. Most probably the availability of the common host, sheep at Krk, may have decreased dramatically in the year of investigation. For its development, this tick species needs one blood donation per year and stage (HOOGSTRAAL et al. 1981), so the ticks may have tried to escape starvation by a 'panic behaviour'. (ii) If this is true, the *Hepatozoon* species found is either not a true reptilian parasite, because this parasite may be passed transstadially by these ticks, but not vertically between different host individuals within one year, or this *Hepatozoon* species is a true reptilian parasite but uses another mite or tick as a vector.

ACKNOWLEDGEMENTS: The author thanks the Austrian Herpetological Society (ÖGH), especially Werner MAYER (Vienna, A), for organisational support, and P. ZWART (Bunnik, NL) for inestimable advice.

REFERENCES: BABOS, S. (1964): Die Zeckenfauna Mitteleuropas; Budapest (Akademiai Kiado), 410 pp. HOOGSTRAAL, H. & WASSEF, H. Y. & BÜTTIKER, W. (1981): Ticks (Acarina) of Saudi Arabia Fam. Argasidae, Ixodidae. -Fauna of Saudi Arabia, Basel; 3: 25-110. JANSEN, M. (2002): Ticks infestation in *Lacerta agilis* (LINNAEUS, 1758) and *Zootoca vivipara* (JACQUIN, 1787) in the Spessart.- Salamandra, Frankfurt/M; 38: 85-94. MALKMUS, R. (1995): Starker Befall einer Bergeidechse (*Lacerta vivipara* JACQUIN, 1787) durch die Zecke *Ixodes ricinus* L. 1758 (Rept.: Sauria; Arachn.: Acari).- Nachr. Naturwiss. Mus., Aschaffenburg; 102: 33-36. MERDIVENCI, A. (1969): Türkiye Keneleri Üzerine Arastirmalar; Istanbul (Kutulumus Matbaasi), 420 pp. NOSEK, J. & LICHARD, M. & SZTANKAY, M. (1967): The ecology of ticks in the Tribec and Hronsky Inovec Mountains.- Bull. World Health Org., Genève; 36 (Suppl.1): 49-59. RATHBAUER, F. (2002): Feldherpetologische Exkursion nach Cres. - ÖGH-Aktuell, Wien, 10: 6-10. REHACEK, J. & NOSEK, J. & GRESKOVA, M. (1961): Study of the relation of the green lizard (*Lacerta viridis* LAUR.) to the natural foci of tick-borne encephalitis.- J. Hygiene, Epidemiol., Mikrobiol. Immunol., Prague; 5: 366-372. SEHNAL, P. (1999): Zur Individuendichte der Adriatischen Mauereidechse, *Podarcis melisellensis fumana* (WERNER, 1891) auf Weideflächen südwestlich von Srem (Cres, Kroatien) (Squamata: Sauria; Lacertidae).- Herpetozoa, Wien, 12: 157-160.

KEYWORDS: Arachnida: Acari: Ixodidae: *Haemaphysalis concinna*, nymph, vector, reptilian hosts (*Algyroides nigropunctatus*, *Podarcis melisellensis*, *P. muralis*, *Lacerta bilineata*, *Elaphe longissima*), island of Krk, Croatia

SUBMITTED: March 11, 2003

AUTHOR: ao Univ.-Prof. Dr. Andreas HASSL, Ameisgasse 63/4/12, A-1140 Vienna, Austria < andreas.hassl@univie.ac.at >

Status and new records of *Dasypeltis scabra* (LINNAEUS, 1758), in Morocco

Dasypeltis scabra (LINNAEUS, 1758) is a colubrid which inhabits subsaharan Africa as far as to the Cape in the South. The populations of the snake in south-western Morocco (BONS & GENIEZ 1996), Egypt (ANDERSON 1898), and south-western Arabia (GASPERETTI 1988) are considered Afro-tropical relicts. The species is extremely rare in Morocco (BONS & GENIEZ 1996; SCHLEICH et al. 1996). While the Egyptian Nile valley is a northward corridor of the Ethiopian region for some subsaharan reptile and amphibian species, e.g., for *D. scabra*, the Moroccan population is isolated from the subsaharan ones. In Morocco, the Common Egg-eating Snake was previously known from six records only located along an axis Agdz-Agadir-Tantan-Bou Kraa. Details were published for all these records (listed in BONS & GENIEZ 1996) except for the most recent one made by Gilles TROCHARD who found a dead juvenile specimen eaten by a scorpion under a stone on April 14, 1993 at the village of Dar Lahoussine (fig. 1). In addition, SCHLEICH et al. (1996) present a picture of a captive Common Egg-eating Snake from Morocco without precise geographical origin.

One of us (MG), together with Karim AMRI, found two more specimens of *D. scabra* in Morocco on March 25, 2002. The first specimen was found dead on the road, 29 km from Sidi Ifni towards Goulimine (= Guelmim) [29°12'49"N / 10°4'0"W], in a quite arid shrubby hilly environment. It is an adult female deposited in the collection of the Laboratoire de Biogéographie et Ecologie des Vertébrés de l'Ecole Pratique des Hautes Études (E.P.H.E.) in Montpellier [BEV.7254] (fig. 2). Its principal measurements and counts are as follows: snout-vent length: 53.5 cm; tail length: 7.3 cm; 25 rows of dorsal scales at midbody; 225 ventral plates (anal excluded); 50 pairs of subcaudal plates. The second specimen, also an adult female, was found alive under a stone 16.5 km from Sidi Ifni towards Tiznit near the coastal road [29°29'30"N / 10°4'30"W] (figs. 3, 4), in quite the same environment as the previous one. Its principal measurements and